The Canadian Field-Naturalist

Note

An update to the distribution of the Endangered False Hop Sedge (Carex lupuliformis Sartwell ex Dewey; Cyperaceae) in Ontario

JESSICA A. CONSIGLIO^{1,*} and MICHAEL J. OLDHAM²

Consiglio, J.A., and M.J. Oldham. 2020. An update to the distribution of the Endangered False Hop Sedge (*Carex lupuliformis* Sartwell ex Dewey; Cyperaceae) in Ontario. Canadian Field-Naturalist 134(2): 171–174. https://doi.org/10.22621/cfn. v134i2.2451

Abstract

We provide an update to the Ontario distribution of False Hop Sedge (*Carex lupuliformis*), prompted by the first documented report of the species in the Niagara region, which represents a noteworthy eastern range extension in the province.

Key words: False Hop Sedge; Carex lupuliformis; Lupulinae; Cyperaceae; Niagara Region; Ontario; endangered species; range extension

False Hop Sedge (Carex lupuliformis Sartwell ex Dewey), which occurs in Canada in the southernmost portions of Ontario and Quebec, is federally listed as Endangered (SARA Registry 2019). In Quebec, False Hop Sedge is provincially listed as Threatened, with 14 extant populations as of 2010 (Environment Canada 2014). The species is also listed as Endangered in Ontario (OMNRF 2017) and was described as one of the province's rarest sedges in the Atlas of the Rare Vascular Plants of Ontario (Argus et al. 1982–1987). In the most recent edition of Rare Vascular Plants of Ontario (Oldham and Brinker 2009), False Hop Sedge was listed with a subnational rank (SRank) of S1, Critically Imperilled, which remains its current provincial rank. In the last decade, this species has been recorded in two new counties in southwestern Ontario (Lambton and Niagara), extending its known range in the province to six counties, including extant and extirpated populations. Neither of these new locations was mapped in the most recent Committee on the Status of Endangered Wildlife in Canada (COSEWIC) status report (COSEWIC 2011: Figure 7). The Niagara location, reported here, is a significant eastward range extension.

Of the genus *Carex*, the section *Lupulinae* comprises six species and is endemic to central and east-

ern North America (Reznicek and Ball 1974). This section is characterized by the largest perigynia of all *Carex* species, as well as inflated perigynia with three stigmas and trigonous (three-sided) achenes (Jones and Hatch 1990). In Ontario, section *Lupulinae* is represented by four species: Gray's Sedge (*Carex grayi* J. Carey), False Hop Sedge, Hop Sedge (*Carex lupulina* Muhlenberg ex Willdenow), and Bladder Sedge (*Carex intumescens* Rudge). Hop Sedge may be challenging to differentiate from False Hop Sedge; however, significant differences in the shape of the achene make this distinction possible, as does the overall larger size of False Hop Sedge. In Ontario, mature achenes are present from early July to late October (Leslie 2018).

The achenes of Hop Sedge are rhomboid, often longer than wide and trigonous with faces flat to slightly concave, enclosed in inflated perigynia and ascending on the culm (Ball and Reznicek 2003). In contrast, the achenes of False Hop Sedge are rhombic, often as long as wide and trigonous with faces strongly concave and angles thickened with prominent nipple-like knobs (Figure 1); they are enclosed in inflated perigynia and held ascending to spreading on the culm (Ball and Reznicek 2003).

The habitat of False Hop Sedge has generally been

A contribution towards the cost of this publication has been provided by the Thomas Manning Memorial Fund of the Ottawa Field-Naturalists' Club.

¹Credit Valley Conservation, 1255 Old Derry Road, Mississauga, Ontario L5N 6R4 Canada

Natural Heritage Information Centre, Ontario Ministry of Natural Resources and Forestry, 300 Water Street, Peterborough, Ontario K9L 3C8 Canada

^{*}Corresponding author: consiglio.jessica@gmail.com



FIGURE 1. Mature False Hop Sedge (*Carex lupuliformis*) achene, from plant material collected in Niagara Region. Photo: Jessica Consiglio.

described as wet forests, where it prefers canopy openings, as well as riverine wetlands, marshes, and wet thickets on calcareous soil (Thompson and Paris 2004). Ontario sites are typically treed swamps with extensive vernal pooling, characterized by Red Maple (Acer rubrum L.), Silver Maple (Acer saccharinum L.), or Green Ash (Fraxinus pennsylvanica Marshall) growing on clay loam (OMNRF 2017). In Quebec, False Hop Sedge is predominantly found growing in canopy openings of Silver Maple swamps (Environment Canada 2014; Langlois and Pellerin 2016). Further south, common associates at sites in Connecticut and Massachusetts include Red Maple, Black Gum (Nyssa sylvatica Marshall), Green Ash, Swamp White Oak (Quercus bicolor Willdenow), and Pin Oak (Quercus palustris Münchhausen) in areas with extensive vernal pooling (Thompson and Paris 2004).

False Hop Sedge was first discovered in Ontario by W. Herriot who collected a specimen in 1902 from Waterloo region, although this population has not been seen again despite several attempts to relocate it (Oldham *et al.* 1993). As reported by Oldham and Crins (1988), A.A. Reznicek, M.J.O., and W. Botham rediscovered the species in Ontario in 1985 at a site near Amherstburg, Essex County, where ~15 plants were growing with Eastern Buttonbush (*Cephalanthus occidentalis* L.) and Hop Sedge in a wet willow floodplain woods. Oldham *et al.* (1993) reported a new population discovered in 1992 growing in a small depres-

sion at the edge of a deciduous woods in the City of London, Middlesex County, and two additional new sites in Aldborough Township, Elgin County, discovered in 1993. At all three sites, False Hop Sedge was observed growing in the open or at the edges of treed areas that held water in the spring (Oldham et al. 1993). The following year, another population was discovered in Middlesex County, near Mount Brydges, by Ron Vanderjeugd (Natural Heritage Information Centre [NHIC] files).

Over a decade later, a new population was documented several kilometres west of West Lorne, Elgin County, by staff from various agencies conducting targetted searches for the species in 2005 (NHIC files). Several years later, Ministry of Natural Resources (MNR) staff documented the species in Ailsa Craig, Middlesex County, growing in a Silver Maple swamp with associated species, such as Blue Beech (Carpinus caroliniana Walter), poison ivy (Toxicodendron Miller spp.), Wild Geranium (Geranium maculatum L.), jewelweed (Impatiens L. spp.), and Fowl Manna Grass (Glyceria striata Lamarck; NHIC files). In 2009, a new population was discovered by Charles Cecile in London, consisting of a few plants growing in an open, moist Silver Maple woodland located between two small wetlands (NHIC files). In addition, MNR staff observed a new False Hop Sedge population in 2011 in Southwold Township, Elgin County, growing in a treed swamp. In 2015, Brian Miller discovered two new False Hop Sedge populations: one growing in woods north of Dresden, Lambton County, which was the first record of this species in the county, and a third new population found in a wooded swamp in London (NHIC files).

The newest recorded Ontario population of False Hop Sedge was discovered in early September 2019 by J.A.C. during a field trip with the Field Botanists of Ontario (Figure 2), led by Albert Garofalo and Nate Torenvliet in Niagara Falls. Approximately 50 clumps were found growing at the edge of extensive vernal pools; associated species included Red Maple, hickory (Carya Nuttall spp.), Black Gum, Pin Oak, and Highbush Blueberry (Vaccinium corymbosum L.). Three additional patches, containing similar numbers of clumps, were subsequently found by Andrew Dean, Nadia Cavallin, and Lisa Riederer during the course of the field trip within a few kilometres of the first patch. These represent the first recorded populations of False Hop Sedge in the Niagara region and a significant eastward range extension for the species in Ontario, as it is ~200 km from the nearest presumed extant population.

With a recent increase in the known range of the species in Ontario, it is worth discussing what may have led to False Hop Sedge being historically over-




FIGURE 2. Range map of all known Ontario False Hop Sedge (*Carex lupuliformis*) populations (extirpated and known populations; COSEWIC 2011: Figure 7). On the far right is the newly discovered population in Niagara. Map: Jessica Consiglio. Basemap sources: Esri, HERE, Garmin, © OpenStreetMap contributors and the GIS user community.

looked. For instance, the visually similar Hop Sedge has been reported growing together with False Hop Sedge at most Ontario sites, which could make detection more challenging if botanists are not aware of the subtle differences mentioned above. However, it is not simply a lack of awareness of the species, as a search of major Ontario herbaria by A.A. Reznicek (1973) during his M.Sc. thesis research on the taxonomy of Carex series Lupulinae in Canada, did not reveal any misidentified material of False Hop Sedge filed with other members of the section (Oldham et al. 1993). This is the case outside Ontario as well, with many North American herbarium records suggesting that False Hop Sedge has never been common or abundant, and populations throughout its range generally consist of ~15 individuals (NatureServe 2019).

Despite possible challenges to detection, or perhaps because of them, field botanists in Ontario and Quebec should remain on the lookout for this rare species, especially from July to October when the mature achenes allow for identification.

Voucher specimens

In accordance with guidance given under the *Endangered Species Act*, 2007 (ESA) the specimen was donated to the Ontario NHIC herbarium.

CANADA, ONTARIO: Lambton County, near Dresden, 10–20 small plants in low depression in woods, 2 September 2015, *M.J. Oldham and B. Miller 43139*

(MICH, NHIC 12029).

CANADA, ONTARIO: Niagara Region, Niagara Falls. Black Gum–Red Maple deciduous swamp, associated with *Acer rubrum*, *Nyssa sylvatica*, *Quercus palustris*, *Vaccinium corymbosum*. 7 September 2019, *J.A. Consiglio* (NHIC 09600).

Acknowledgements

Thanks to Nadia Cavallin of the Royal Botanical Gardens (Hamilton) for providing access to the camera equipment used for the achene photo and to Corey Burt for his assistance with the camera software. Eric Snyder provided helpful guidance on Ontario Endangered Species Act considerations. Many thanks to Dawn Renfrew for her input on an early draft and to reviewers Pauline Catling and Tony Reznicek, whose thoughtful comments improved the quality of this note. The insightful feedback provided by Associate Editor, Paul Catling, was also beneficial and much appreciated.

Literature Cited

Argus, G.W., K.M. Pryer, D.J. White, and C.J. Keddy. 1982–1987. Atlas of the Rare Vascular Plants of Ontario. 4 parts. National Museum of Natural Sciences, Ottawa, Ontario, Canada.

Ball, P.W., and A.A. Reznicek. 2003. Carex. Pages 252–258 in Flora of North America North of Mexico (Vol. 23). Edited by Flora of North America Editorial Committee.

- Flora of North America Editorial Committee, New York, USA, and Oxford, United Kingdom. Accessed 5 June 2020. http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=105644.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2011. COSEWIC assessment and status report on the False Hop Sedge *Carex lupuliformis* in Canada. COSEWIC, Ottawa, Ontario, Canada.
- Environment Canada. 2014. Action plan for the False Hop Sedge (*Carex lupuliformis*) in Quebec. *Species at Risk* Act action plan series. Environment Canada, Ottawa, Ontario, Canada. Accessed 9 May 2020. https://tinyurl. com/yaol5gjy.
- Jones, S.D., and S.L. Hatch. 1990. Synopsis of Carex section Lupulinae (Cyperaceae) in Texas. Sida 14: 87–99.
- Langlois, A., and S. Pellerin. 2016. Recovery of the endangered False Hop Sedge: a ten-year study. Global Ecology and Conservation 7: 214–224. https://doi.org/10.1016/j.gecco.2016.07.003
- Leslie, J. 2018. Vascular Plants at Risk in Ontario. Accessed 7 January 2019. https://tinyurl.com/y6quh2jc.
- NatureServe. 2019. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia, USA. Accessed 18 February 2020. http://explorer.natureserve.org.
- Oldham, M.J., and S.R. Brinker. 2009. Rare vascular plants of Ontario. Fourth edition. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada. Accessed 17 December 2019. https://tinyurl.com/y4v7chjm.
- Oldham, M.J., and W.J. Crins. 1988. New and significant records of Ontario sedges (Cyperaceae). Canadian Field-

- Naturalist 102: 500–507. Accessed 17 December 2019. https://www.biodiversitylibrary.org/page/28243320.
- Oldham, M.J., D. McLeod, and R. Vanderjeugd. 1993.
 Carex lupuliformis (Cyperaceae) new to Elgin and Middlesex Counties, Ontario. Field Botanists of Ontario Newsletter 6(3): 17–18.
- OMNRF (Ontario Ministry of Natural Resources and Forestry). 2017. Recovery strategy for the False Hop Sedge (*Carex lupuliformis*) in Ontario. Ontario recovery strategy series. Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario, Canada. Accessed 17 December 2019. https://www.ontario.ca/page/false-hop-sedge-recovery-strategy.
- Reznicek, A.A. 1973. The taxonomy of *Carex* series *Lupulinae* in Canada. M.Sc. thesis, University of Toronto, Toronto, Ontario, Canada.
- Reznicek, A.A., and P.W. Ball. 1974. The taxonomy of Carex series Lupulinae in Canada. Canadian Journal of Botany 52: 2387–2399. https://doi.org/10.1139/b74-310
- SARA (*Species at Risk Act*) Registry. 2019. Species profile: False Hop Sedge (*Carex lupuliformis*). Accessed 5 June 2020. https://species-registry.canada.ca/index-en. html#/species/2-13.
- Thompson, E., and C.A. Paris. 2004. Carex lupuliformis Sartwell ex Dewey: False Hop Sedge. Conservation and research plan for New England. New England Wild Flower Society, Framingham, Massachusetts, USA. Accessed 17 December 2019. https://tinyurl.com/y9y7c9tt.

Received 26 February 2020 Accepted 5 June 2020 Associate Editor: P.M. Catling