

The freshwater mussel (Bivalvia: Unionidae) assemblage of the Mississagi River, Ontario: a stronghold for the federally Endangered Hickorynut (*Obovaria olivaria*)

KELLY A. McNICHOLS-O'ROURKE^{1,*}, MANDY P. GIBSON¹, KEITH SAYERS², and TODD J. MORRIS¹

¹Fisheries and Oceans Canada, Great Lakes Laboratory for Fisheries and Aquatic Sciences, 867 Lakeshore Road, Burlington, Ontario L7S 1A1 Canada

²Mississauga First Nation Lands and Resources Department, 64 Park Road, P.O. Box 1299 Mississauga First Nation, Blind River, Ontario P0R 1B0 Canada

*Corresponding author: kelly.mcnichols-orourke@dfo-mpo.gc.ca

McNichols-O'Rourke, K.A., M.P. Gibson, K. Sayers, and T.J. Morris. 2026. The freshwater mussel (Bivalvia: Unionidae) assemblage of the Mississagi River, Ontario: a stronghold for the federally Endangered Hickorynut (*Obovaria olivaria*). Canadian Field-Naturalist 139(2): 85–100. <https://doi.org/10.22621/cfn.v139i2.3439>

Abstract

Federally Endangered Hickorynut (*Obovaria olivaria*) was historically known to occupy a short section of the Mississagi River. Whether this distribution represented the true range of the species or simply reflected a paucity of directed mussel sampling was unclear. The Mississagi River was systematically surveyed in 2023 to evaluate the freshwater mussel assemblage in the watershed, focussing on Hickorynut. The river was divided into 21 contiguous reaches, the first of which was 1 km long, followed by 20, 2 km reaches from the Red Rock Falls Generating Station downstream to the mouth at Lake Huron. Each reach was surveyed at least once using SCUBA; additional surveys were conducted in some reaches if warranted (e.g., historical presence of Hickorynut). In total, 35 sites were surveyed (26 SCUBA, nine snorkel) and 7590 mussels of seven species, including 141 Hickorynut were found. A single Lake Floater (*Pyganodon lacustris*) was found and genetically verified, representing the third confirmed detection of the species in Canada. Eastern Elliptio (*Elliptio complanata*) was the most common species found, occurring at all sites. Our results clarify the distribution of mussels throughout this stretch of the Mississagi River and extend the known distribution of Hickorynut ~22 km upstream of its previously known range. These findings will aid in the conservation and management of freshwater mussels in Canada and inform future assessments of at-risk mussel species.

Key words: Unionidae; Mollusc; Species-at-Risk; Lake Huron

Introduction

Freshwater mussels (Unionidae) are widely distributed animals found in lakes, rivers, canals, wetlands, and agricultural drains where they can represent the largest components of the benthic biomass and are important contributors to aquatic ecosystems (Vaughn and Hoellein 2018). Vaughn (2018) describes in detail important ecosystem services mussels provide (e.g., biofiltration, nutrient cycling and storage, environmental monitoring, food, culture) that benefit aquatic ecosystems and society in general. Unfortunately, these important animals are globally imperilled with declines recorded (summarized in Aldridge *et al.* 2022) because of pollution, habitat degradation and modification, water quality, invasive species, over-harvest, climate change, and loss of host fishes (Nobles and Zhang 2011; Haag and Williams 2014; Lopes-Lima *et al.* 2018; Ferreira-Rodríguez *et*

al. 2019; Hayes Morris and Morris 2024). Approximately 67% of North American freshwater mussel species are considered threatened or near-threatened (Lopes-Lima *et al.* 2018). The Committee on the Status of Endangered Wildlife in Canada (COSEWIC) has assessed 37% of the 55 species of unionids in Canada as at-risk (Extirpated, Endangered, Threatened, or Special Concern; Government of Canada 2023).

Hickorynut (*Obovaria olivaria*), was assessed as Endangered by COSEWIC in 2011 and listed as such under the provincial *Endangered Species Act* in 2012 (Government of Ontario 2014) and under the Federal *Species at Risk Act* in 2019 (SARA Public Registry 2024a). The main threats were identified as the direct and indirect effects of invasive dreissenid mussels and reduced abundance of their host fish (COSEWIC 2011). Like all Unionidae freshwater mussels in Canada, Hickorynut has a complex lifecycle that requires

a period of glochidial encystment on a vertebrate host. The suspected host for Hickorynut in Canada is Lake Sturgeon (*Acipenser fulvescens*; Brady *et al.* 2004). The Great Lakes – Upper St. Lawrence population of Lake Sturgeon in the range of Hickorynut in Canada is also considered at-risk and was assessed by COSEWIC as Threatened in 2006, a status reconfirmed in 2017 (SARA Public Registry 2024b). Major threats include harvest, dam creation/operation (habitat alteration, barriers, entrainment losses), invasive species, and pollution (COSEWIC 2017).

Across its North American distribution, Hickorynut status ranges from “apparently secure” to “presumed extirpated” (NatureServe 2026). Hickorynut was historically reported from the watersheds of Lake Huron, Lake Erie, Lake Ontario, and the St. Lawrence River; however, it was never found in the Canadian portion of the Lake Superior watershed (Bell 1861; Metcalfe-Smith *et al.* 2000; Martel *et al.* 2006; Schloesser *et al.* 2006; COSEWIC 2011; Martel and Madill 2017; Martel *et al.* 2023; LGLUD 2024). Currently, within Canada, Hickorynut is extant only in the Mississagi River (Lake Huron drainage) and the St. Lawrence and Ottawa river systems (Martel and Picard 2005; Martel *et al.* 2006, 2023; COSEWIC 2011; Martel and Madill 2017), with the St. Lawrence River and its tributaries having the largest remaining Canadian populations.

Hickorynut was first collected from the Mississagi River in 1859–1860 by Dr. R. Bell (Bell 1861; Table 1) with no additional reports until almost a century later in 1955, and then again in 2000 (COSEWIC 2011; Zanatta and Woolnough 2011; Harvard University Museum of Comparative Zoology 2016). While

preparing the 2011 COSEWIC assessment, Zanatta and Woolnough (2011) undertook a small survey and confirmed the presence of an extant and reproducing population in 2009 in the Mississagi River, with additional Hickorynut found in 2017 (Buchholz *et al.* 2022). Following the 2011 COSEWIC assessment, Fisheries and Oceans Canada (DFO) conducted a recovery potential assessment on Hickorynut in Canada in 2013 and rated the status of the Mississagi River population as “poor” because of its low relative abundance index and unknown population trajectory (Bouvier *et al.* 2013). Given the limited information available on Hickorynut in the Mississagi River, we undertook an assessment of the entire freshwater mussel assemblage of the river in 2023. The goal was to determine freshwater mussel distribution, with a focus on the Endangered Hickorynut, throughout the Mississagi River from the Red Rock Generating Station to the mouth of the river along the north shore of Lake Huron. The river represents an isolated population of Hickorynut that is the only extant population of the species upstream of the St. Lawrence River. Therefore, a thorough understanding of the distribution of this species as well as the entire mussel assemblage will aid in future conservation and management efforts.

Methods

The Mississagi River watershed is on the north shore of Lake Huron adjacent to the Mississauga First Nation (Figure 1). The river is predominantly within the Sudbury and Algoma districts (Hatch 2010, 2022). It drains an area of ~9300 km², originates upstream of White Owl Lake and empties into a delta in the North

TABLE 1. Freshwater mussel species found in the Mississagi River, Ontario, over time; S indicates a shell, not a live individual. Nomenclature from Bell (1861) has been updated and scientific names follow MolluscaBase (2023). The single mussel species-at-risk, Hickorynut (*Obovaria olivaria*), is highlighted in grey.

Scientific name	Common name	1859–1860 (Bell 1861)	2009 (Zanatta and Woolnough 2011)	2023 (our survey)
<i>Elliptio complanata</i>	Eastern Elliptio	X	X	X
<i>Lampsilis cardium</i>	Plain Pocketbook	X	X	X
<i>Lampsilis siliquoidea</i> *	Fatmucket		X	X
<i>Lampsilis radiata</i> *	Eastern Lampmussel	X		
<i>Lasnigona costata</i>	Flutedshell	X		X (S)
<i>Ligumia recta</i>	Black Sandshell	X	X (S)	X (S)
<i>Alasmidonta marginata</i>	Elktoe	X		
<i>Obovaria olivaria</i>	Hickorynut	X	X	X
<i>Pyganodon grandis</i>	Giant Floater		X	X
<i>Pyganodon lacustris</i>	Lake Floater			X
<i>Strophitus undulatus</i>	Creeper			X

*These two species can be difficult to differentiate morphologically, are known to hybridize (Porto-Hannes *et al.* 2021), and the taxonomy has historically been fluid (MolluscaBase 2025).




FIGURE 1. Mussel abundance (a) and species richness (b) at each site sampled in the Mississagi River, Ontario, in 2023.

Channel of Lake Huron west of Blind River, Ontario (Hatch 2010). The watershed is sparsely populated by humans and characterized by large lakes and wetlands (Hatch 2022); parts of it flow through several provincial parks (MECP 2024). The entire river contains a diverse fish assemblage because it has a variety of habitats (CNSC 2023; DFO's Aquatic Invasive Species Program unpubl. data). There are four water-power generating stations along the Mississagi River (Hatch 2010) and our study focusses on the ~30 km stretch that begins downstream of the Red Rock Falls Generating Station (the first dam upstream of Lake Huron) and continues to the river mouth. This Generating Station is a functional barrier to host fish passage (Barth *et al.* 2018) and was used as the upstream boundary for our survey. Substrate in this area largely consists of sand and silts with some small mix of gravel and cobble (Ontario Hydro 1982, as cited in Hatch 2010). The delta at the mouth of the river is considered provincially significant given its natural and cultural features (e.g., "bird's foot" delta formation, plant communities, and breeding and migrating birds) and is part of the Mississagi Delta Provincial Park and Nature Reserve—protecting an area of over 2300 ha (MECP 2021).

We systematically divided the river into 21 reaches beginning downstream of the Red Rock Generating Station and extending to the mouth of the river using Google My Maps (Figure 1) to more evenly allocate effort and to ensure coverage of the entire river. The upper most section was only 1 km long given the proximity to the Generating Station and higher water velocity in this area. All other reaches were 2 km long. The portion of the river extending through the delta contains numerous individual channels, therefore using the largest channels, we divided the delta into three main sections with each beginning at the site downstream of the Mississagi River Falls (site 23 in Figure 1). Each section included two reaches that were sampled and when the reaches extended into Lake Huron, sample sites were restricted to areas in the river. Within each reach, the sample site for a SCUBA survey was selected using four criteria: historical presence of Hickorynut (three of the four known occurrence sites were resurveyed), suitable substrate (some evidence of sand substrate), suitable depth (2–5 m), and diver safety (i.e., no visible hazards, safe flow and current, low boat traffic). A single site, where a fresh Hickorynut shell was recorded in 2000 by M.J. Oldham, was not resurveyed in 2023 because of safety concerns (i.e., its proximity to the Mississagi River Falls).

In addition to the single SCUBA site in each reach, some reaches received additional survey effort either via SCUBA or snorkelling. Additional SCUBA sites

were conducted in reaches where no evidence of Hickorynut occurred during the first survey or if only shells were found within a given reach and additional time was available. Snorkelling sites were placed in shallow areas of reaches where Hickorynut was collected during the initial SCUBA surveys, had an appropriate snorkel depth (≤ 2 m depth), and contained preferred Hickorynut habitat. While snorkelling is a less expensive and safer method to assess potential Hickorynut habitat in shallower areas, it was expected that SCUBA would be required and be more successful in deeper water because of the mussel's depth preference.

Surveys were conducted using methods similar to Keretz *et al.* (2021). Two vessels were present at each SCUBA site, one dive boat and one processing boat, whereas one vessel was present at snorkel sites. At SCUBA sites, the dive boat team was tasked with collecting mussels via SCUBA and assessing underwater site conditions, whereas the processing boat team was responsible for the collection of above/surface-water site data and mussel processing. Each site was surveyed visually or tactilely by either two SCUBA divers or two snorkellers for 30 min. If >10 individual mussels, three species, or any mussel species-at-risk were detected, sampling continued for an additional 10 min by each diver/snorkeller, provided environmental conditions (e.g., water velocity) were suitable. SCUBA divers/snorkellers moved randomly around the vessel or along the shoreline, collecting unionid mussels. There was no overlap in the areas searched by each diver/snorkeller. When large patches and high densities of Eastern Elliptio (*Elliptio complanata*) were observed, divers/snorkellers collected some individuals and counted the others to save processing time and prevent unnecessary handling of the animals. All mussels found were identified to species, sexed visually (dimorphic species only), and counted. A maximum of 20 individuals of each common species (a variety of sizes were selected) and all Hickorynut were measured (maximum length) using Vernier calipers. Shells or valves found were also identified to species. After processing, all live mussels were returned to the site from which they were collected, unmarked. For all individuals that were identified to species using shell morphology, vouchers in the form of either photographs and/or shells were collected (Morris *et al.* 2022; Figures S1–S9). If positive species identification could not be made in the field, a sample was collected for genetic identification. For small or thin-shelled specimens the whole animal was collected following Millar *et al.* (2022); a non-destructive sample was taken of larger specimens using a Cytology brush, where the viscera/foot was swabbed following Henley *et al.* (2006) and Harrison *et al.* (2023).

Molecular analyses were completed by the Hanner

Lab at the University of Guelph. Foot tissue from each mussel was used for lysis and DNA extraction with a Qiagen DNeasy Blood & Tissue Kit (Hilden, North Rhine-Westphalia, Germany). For polymerase chain reaction (PCR) and sequencing, Folmer *et al.* (1994) primers in a standard 25 μ L PCR reaction were used. The PCR products were screened on a gel and purified with magnetic beads. Bidirectional Sanger sequencing was performed at the Genomics Facility, Advanced Analysis Centre at the University of Guelph. Sequences were edited and aligned in Geneious Prime (<https://www.geneious.com>) and analyzed in the Barcode of Life Data Systems (BOLD; Ratnasingham and Hebert 2007).

Site-level environmental data were collected at each site and included: air temperature using a Kestrel 2000 Pocket Wind Meter (Nielsen-Kellerman Co., Boothwyn, Pennsylvania, USA); water temperature ($^{\circ}$ C), conductivity (μ S/cm), total dissolved solids (TDS, mg/L), salinity (psu), dissolved oxygen (optical dissolved oxygen, ODO, % and mg/L), pH, and turbidity (FNU) using a YSI EXO2 multiparameter sonde (YSI Inc., Yellow Springs, Ohio, USA); surface (top 10 cm) water velocity (m/s) using an OTT MF Pro flow meter (OTT HydroMet, Kempten, Germany); and, water clarity (m) using a 1.2 m Secchi tube. The minimum, maximum, and mean width of the river at each site was measured using a Nikon Laser 1200S waterproof laser range finder (Nikon Canada Inc., Mississauga, Ontario, Canada). Maximum and mean depths searched were measured by SCUBA divers using either Shearwater Perdix 2 or Shearwater Nerd technical dive computers (Shearwater Research Inc., Richmond, British Columbia, Canada). Substrate composition (%) was visually estimated by divers using Stanfield (2010) substrate class categories. Box plots at sites with (live) and without (no live or shells) live Hickorynut were created for all environmental data collected at the site level using Microsoft Excel for Microsoft 365. In addition, a principal component analysis (PCA) was performed in R (version 4.3.0; R Core Team 2023) using the package “vegan” (Oksanen *et al.* 2022) to evaluate the relationship between substrate and Hickorynut presence/absence. Prior to analysis, substrate data were transformed using arcsine transformation. All non-substrate habitat parameters were excluded from the PCA because of limited data, and correlation with other variables.

Catch-per-unit-effort (CPUE) was calculated by comparing the number of mussels found to the amount of time searched:

$$\text{CPUE} = \frac{\sum M_i}{\#PH}$$

where $\sum M_i$ is the sum of all mussels found (either all individuals or individuals of a specific species)

at each site sampled and #PH is the total number of person-hours (PH) searched (e.g., two people each searching for 0.5 h is equal to one PH of search time).

Results

Thirty-five sites, totalling 44.93 PH of search effort, were surveyed in the Mississagi River between 25 and 30 July 2023 (Tables 2 and 3). Each of the 21 reaches was sampled from one to three times (Figure 1). Twenty-six sites were sampled using SCUBA, including three sites previously known to have Hickorynut, and nine were sampled using snorkel (Table 2). All sites, except for three, were sampled for 1.33 PH. The river bottom current at sites 1, 3, and 12 was strong and therefore, as a safety precaution, the SCUBA divers could only search for 1.00, 0.70, and 0.67 PH, respectively.

A total of 7590 mussels, representing seven extant species were observed (Table 3; Figures S1–S7). An additional two species, Flutedshell (*Lasmigona costata*) and Black Sandshell (*Ligumia recta*), represented by shells only were also observed (Table 3; Figures S8–S9). Abundance varied from one to 1143 animals/site (Table 3, Figure 1a); species richness ranged from one to six (Figure 1b). Mean CPUE for all species was $167.48 \pm \text{SE } 27.40$ mussels/PH and ranged from 1.00 to 859.40 mussels/PH (Table 3).

The most widespread and abundant species was Eastern Elliptio, which was found at all sites ($n = 35$) and made up 88% of the individuals observed ($n = 6693$; Table 3). The CPUE for this species ranged from 1.00 to 765.41 mussels/PH (Table 3) with a mean of 147.31 ± 26.68 . Lengths of 647 individuals ranged from 15.30 mm to 100.50 mm (Figure 2).

We were unable to positively identify 26 individuals in the field and they were sent for genetic identification. Sixteen individuals had identifications confirmed via molecular analysis. These included eight Fatmucket (*Lampsilis siliquoidea*), three Giant Floater (*Pyganodon grandis*), three Eastern Elliptio, one Plain Pocketbook, and one Lake Floater. Photographs of the remaining 10 individuals were studied by experts and five could be identified to species: one Creeper (site 18), one Giant Floater (site 32), one Eastern Elliptio (site 23), and two Fatmucket (site 32). Through molecular analysis and shell morphology, two species were detected alive for the first time in the Mississagi River—Lake Floater (*Pyganodon lacustris*) and Creeper (*Strophitus undulatus*; Table 3), respectively. Of the five remaining individuals that could not be identified to species, two were assigned to genus via photographs: *Lampsilis* sp. (site 24) and *Pyganodon* sp. (site 8). Three individuals remained unknown (sites 17 and 35; Table 3). Voucher specimens of Flutedshell (CMN catalogue ID: CMNML

TABLE 2. Sites sampled in the Mississagi River, Ontario, using SCUBA and snorkelling in 2023. Historical (H) sites were surveyed prior to and in 2023. * represent sites sampled via snorkel. Sites are in upstream to downstream order.

Site code	Latitude	Longitude	Date sampled	Site code	Latitude	Longitude	Date sampled
1	46.3093°N	83.2818°W	28 July	19*	46.2052°N	83.0754°W	29 July
2	46.2998°N	83.2605°W	26 July	20	46.2070°N	83.0638°W	27 July
3	46.2925°N	83.2539°W	28 July	21	46.2044°N	83.0475°W	27 July
4	46.2908°N	83.2423°W	26 July	22 (H)	46.2061°N	83.0250°W	27 July
5	46.2805°N	83.2235°W	26 July	23 (H)	46.1990°N	83.0263°W	25 July
6 (H)	46.2728°N	83.2270°W	26 July	24	46.1929°N	83.0293°W	25 July
7	46.2684°N	83.2137°W	26 July	25	46.1867°N	83.0387°W	25 July
8	46.2581°N	83.2016°W	26 July	26	46.1765°N	83.0505°W	25 July
9*	46.2572°N	83.2014°W	29 July	27	46.1750°N	83.0625°W	28 July
10	46.2510°N	83.1664°W	26 July	28	46.1739°N	83.0470°W	28 July
11	46.2442°N	83.1544°W	27 July	29	46.1873°N	83.0315°W	25 July
12	46.2378°N	83.1454°W	28 July	30*	46.1849°N	83.0309°W	30 July
13*	46.2327°N	83.1335°W	29 July	31*	46.1803°N	83.0319°W	30 July
14	46.2326°N	83.1297°W	27 July	32	46.1700°N	83.0422°W	28 July
15*	46.2263°N	83.1131°W	29 July	33*	46.1817°N	83.0267°W	30 July
16	46.2221°N	83.1094°W	27 July	34*	46.1782°N	83.0194°W	30 July
17*	46.2139°N	83.1019°W	29 July	35	46.1769°N	83.0199°W	25 July
18	46.2094°N	83.0868°W	27 July				

2025-1318, shell; Figure S8), Black Sandshell (CMNML 2025-1317, valve; Figure S9 and CMNML 2025-1319, shell), Hickorynut (CMNML 2025-1316, shell, CMNML 2025-1320, shell and valve), and Lake Floater (CMNML 2025-1321, wet; Figure S6) are stored at the Canadian Museum of Nature.

A total of 141 live Hickorynut were found at 18 sites, 14 via SCUBA and four via snorkel (Figure 3). It was the third most abundant species and the only federal or provincially listed mussel observed in the Mississagi River. It made up <2% of the individuals found, but occurred at ~51% of sites, including a site upstream of the town of Iron Bridge. Shells were observed at an additional six sites increasing the frequency of occurrence to ~69% (Table 3). Mean CPUE for Hickorynut at occupied sites was 5.75 ± 1.99 and ranged from 0.75 to 30.08 mussels/PH (Table 3). Lengths ranged from 32.30 mm to 84.00 mm (Figure 2).

Mean (\pm SE) air and water temperature of all sites sampled were $25.33 \pm 0.47^\circ\text{C}$ and $22.32 \pm 0.19^\circ\text{C}$, respectively (Table S1). Mean YSI EXO2 multiparameter sonde water quality measurements for conductivity, total dissolved solids (TDS), salinity, and pH were $51.13 \pm 0.44 \mu\text{S/cm}$, $35.01 \pm 0.21 \text{ mg/L}$, $0.02 \pm 0.00 \text{ psu}$, and 7.41 ± 0.03 , respectively (Table S1). Additional measurements from the YSI included dissolved oxygen (% and mg/L) and turbidity which were $99.89 \pm 0.87\%$, $8.69 \pm 0.08 \text{ mg/L}$, and $0.63 \pm 0.06 \text{ FNU}$, respectively (Table S1). Mean surface water velocity was $0.24 \pm 0.03 \text{ m/s}$ and water clarity

was over 1.2 m at all but one site (Table S1). The width of the river at sites sampled ranged from 29 to 229.50 m with a mean of 113.88 ± 9.21 (Table S1). The maximum depth surveyed was 16.00 m at site 8 and across all sites the mean depth searched was $2.64 \pm 0.33 \text{ m}$ (Table S1).

As expected, given the close proximity of all sample sites, physical habitat data collected at each site were similar whether or not live Hickorynut were present (Figures S10). Fine sediments (sand, silt, clay, and muck) made up over 75% of the substrates across the 35 sites sampled. The dominant substrate was sand with a mean of $60.71 \pm 4.02\%$ and median of 65% (Table S1). Sites with Hickorynut tended to have higher proportions of sand (Figure 4) whereas sites absent of Hickorynut tended to have higher amounts of coarse material (boulder and cobble) and silt (Figure 4). Similarly, the first two axes of the PCA (explaining over 50% of the observed variation) showed a high degree of overlap among sites with (present) and without (absent) Hickorynut; however, Hickorynut presence was associated with a higher proportion of sand and lower proportion of boulder and cobble (Figure 5).

Discussion

The collaborative effort conducted in 2023 by DFO, Mississauga First Nation Lands Resources Department, and Environment and Climate Change Canada lead to the most extensive and systematic

TABLE 3. Mussel species found at each site surveyed in the Mississagi River, Ontario, in 2023. The single mussel species-at-risk, Hickorynut (*Obovaria olivaria*) is highlighted in grey. S(#) represents species found as complete shells and the number of shells found. V(#) represents species found as valves (one half of a complete shell) and the number of valves found. SH† indicates fresh shell/valve and all others were weathered in condition. Unknown individuals are included in the abundance total, but not in the species richness totals. Effort is represented by the total number (#) of person-hours (PH) surveyed. Catch-per-unit-effort (CPUE) was calculated for all mussels found at a site and for the Endangered Hickorynut and rounded to the nearest whole number. Nomenclature follows MolluscaBase (2023). Continued on the following page.

Scientific name	Common name	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
<i>Elliptio complanata</i>	Eastern Elliptio	1	285	55*	428	41	534	95	155	45	346	109	116	360	113	497	66	90	1018
<i>Lampsilis cardium</i>	Plain Pocketbook	0	0	0	0	0	0	0	1	0	2	0	S(1)	0	3	1	0	1	7
<i>Lampsilis siliquoides</i>	Fatmucket	0	10	12	13	30	37*	9*	23	1	16	18	32	7	80*	0	20	15	75*
<i>Lampsilis</i> sp.		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Lasmigona costata</i>	Flutedshell	0	S(1)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ligumia recta</i>	Black Sandshell	0	0	0	0	0	0	V(3)	0	0	V(1)	0	0	0	V(1)	0	S(2)	0	0
<i>Obovaria olivaria</i>	Hickorynut	0	S(2) V(5)	0	0	1	0	V(2)	1	0	S(1) V(1)	0	0	0	1	0	1	0	40
<i>Pyganodon grandis</i>	Giant Floater	0	0	0	2*	5*	1	0	0	1	0	0	0	V(1)	0	0	0	1	2
<i>Pyganodon lacustris</i>	Lake Floater	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Pyganodon</i> sp.		0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
<i>Strophitus undulatus</i>	Creeper	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1
Unknown		0	0	0	0	V(1)	0	0	0	0	0	0	0	0	0	0	0	0	2
Total abundance		1	295	67	443	78	572	104	181	47	364	127	148	367	197	498	87	110	1143
Live species richness		1	2	2	3	5	3	2	4	3	3	2	2	2	4	2	3	5	6
Total species richness		1	4	2	3	5	3	4	4	3	5	2	3	3	5	2	4	6	6
Effort (# PH)		1.00	1.33	0.70	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	0.67	1.33	1.33	1.33	1.33	1.33	1.33
CPUE all mussels (# mussels/PH)		1.00	221.80	95.71	333.08	58.65	430.08	78.20	136.09	35.34	273.68	95.49	220.90	275.94	148.12	374.44	65.41	82.71	859.40
CPUE <i>Obovaria Olivaria</i> (# mussels/PH)		0.00	S,V	0.00	0.00	0.75	0.00	V	0.75	0.00	S,V	0.00	0.00	0.00	0.75	0.00	0.75	0.00	S

*Species identification confirmed genetically (see Methods).

TABLE 3. Continued.

Scientific name	Common name	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	Total
<i>Elliptio complanata</i>	Eastern Elliptio	406	195	64	128	205	84	151	63	28	35	166	160	37	45	87	122	363	6693
<i>Lampsilis cardium</i>	Plain Pocketbook	0	10	0	1	1	0	0	0	0	1	1	5	2	0	2	5	6*	49
<i>Lampsilis siliquoidea</i>	Fatmucket	7	24	25	2	10	3*	5	2	7	13	7	3	4	109	0	6	24*	649
<i>Lampsilis</i> sp.		0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1
<i>Lasmigona costata</i>	Flutedshell	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	S(1)
<i>Ligumia recta</i>	Black Sandshell	0	V(5)	V(1)	V(1)	0	0	0	0	0	0	V(1)	0	0	0	0	0	0	S(2) V(13)
<i>Obovaria olivaria</i>	Hickorynut	0	22	1	2	19	2	2	SH(1)†	1	S(3) V(3)	4	19	1	0	2	2	20	141
<i>Pyganodon grandis</i>	Giant Floater	0	1	0	0	2	0	3	1	5	4	2	2	3	6	2	0	0	43
<i>Pyganodon lacustris</i>	Lake Floater	0	0	0	0	0	0	0	0	0	0	0	0	0	1*	0	0	0	1
<i>Pyganodon</i> sp.		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
<i>Strophitus undulatus</i>	Creeper	0	0	0	0	1	0	0	0	0	1	1	0	0	0	2	0	1	9
Unknown	Unknown	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	3
Total abundance		413	252	90	133	238	90	161	66	41	54	181	189	47	161	95	135	415	7590
Live species richness		2	5	3	4	6	3	4	3	4	5	6	5	5	4	5	4	5	7
Total species richness		2	6	4	5	6	3	4	4	4	6	7	5	5	4	5	4	5	9
Effort (# PH)		1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33	1.33
CPUE all mussels (# mussels/PH)		310.53	189.47	67.67	100.00	178.95	67.67	121.05	49.62	30.83	40.60	136.09	142.11	35.34	121.05	71.43	101.50	312.03	
CPUE <i>Obovaria Olivaria</i> (# mussels/PH)		0.00	16.54	0.75	1.50	14.29	1.50	1.50	SH	0.75	S,V	3.01	14.29	0.75	0.00	1.50	1.50	15.04	

*Species identification confirmed genetically (see Methods).




FIGURE 2. Length distribution of Eastern Elliptio (*Elliptio complanata*; n = 647) and Hickorynut (*Obovaria olivaria*; n = 141) found in the Mississagi River, Ontario, during 2023 surveys.




FIGURE 3. Hickorynut (*Obovaria olivaria*) presence at sites sampled using SCUBA (circle) or snorkel (triangle) in the Mississagi River, Ontario in 2023.

freshwater mussel survey in the Mississagi River. Mussels were generally abundant, except for the most upstream site where only a single Eastern Elliptio

was found. Eastern Elliptio and Fatmucket were the most abundant and widespread species observed with the federally Endangered Hickorynut the third most




FIGURE 4. Visual estimates of substrate recorded at sites with (present) and without (absent) Hickorynut (*Obovaria olivaria*; \pm standard error). Sampling occurred in the Mississagi River, Ontario in 2023.

abundant species. It is now apparent from our study that the Mississagi River houses a freshwater mussel assemblage of moderate richness with nine species observed, seven extant and two as shells, when compared to other tributaries in the lower portion of Lake Huron (e.g., Saugeen, Maitland, Bayfield,

Assable Rivers) where 14–26 species have been observed (McNichols-O’Rourke *et al.* 2012; Morris *et al.* 2012; DFO 2020; Gibson *et al.* 2024). We know of no standard for richness, although suggest a low species richness in Canada is one or two; 85% of tertiary watersheds in Canada with mussels have fewer




FIGURE 5. Principal component analysis evaluating the relationship between substrate and Hickorynut (*Obovaria olivaria*) presence/absence with ellipses representing the 95% CI of one SD. Sites were sampled in the Mississagi River, Ontario in 2023.

than nine species while 40% have only one or two (DFO unpubl. data).

Our surveys produced new watershed records for two species: Lake Floater confirmed via genetic analysis and Creeper confirmed via shell morphology. Lake Floater was “rediscovered” in Canada in 2018 in Little Sheguiandah Lake, Killarney Provincial Park, Ontario and confirmed through molecular analysis (Morris *et al.* 2020). The species’ distribution has been suggested to comprise “... the St Lawrence, upper Susquehanna and Hudson, upper Mississippi and southern Hudson Bay drainages of North America” (Hoeh and Burch 1989: 269); however, given the difficulty of differentiating species within the *Pyganodon* genus morphologically, Lake Floater has not been broadly identified in Canada and more specimens are required to determine the full Canadian range. The detection of the species in the Mississagi River represents only the third genetically confirmed occurrence in Canada. With only a single Lake Floater individual found, nothing can be said about its reproduction and recruitment without additional sampling. The detection of Creeper in the Mississagi River is not surprising given its known distribution throughout the north shore of Lake Huron (Clarke 1981) and the fact that it is a host generalist (Cliff *et al.* 2001) with many potential host fishes present in the river (Marson *et al.* 2016; DFO’s Aquatic Invasive Species Program unpubl. data). Interestingly, the single Flutedshell observed in 2023 is the only record of the species in the Mississagi River since Bell (1861). These observations highlight our poor understanding of species distributions, even common species, resulting

from a lack of basic surveys in many Canadian rivers.

Based on the abundance and length of individuals observed in most species where more than a single individual was found, it appears that reproduction and recruitment is occurring in the Mississagi River for most of the mussel assemblage. Although several small individuals were found during the surveys, it is important to note that visual surveys have an inherent bias; larger individuals are collected more often than smaller ones (Obermeyer 1998). Small mussels/juveniles are typically burrowed in the sediment and are, therefore, more difficult to find during visual surveys (Miller and Payne 1993; Reid and Morris 2017). It is therefore likely that actual reproduction and recruitment levels are much greater than we detected.

The Mississagi River population of Hickorynut represents the last known population of this species on the Canadian side of the Great Lakes drainage upstream of the St. Lawrence River. It is also the only population known, in Canada or the USA, from the Lake Huron drainage. Given its apparent distributional disjunction, this population represents an important and irreplaceable ecological component of the evolutionary history of this species (Zanatta and Woolnough 2011). Recent genetic work by Bucholz *et al.* (2022), found evidence that Great Lakes Hickorynut populations (including the Mississagi River) are differentiated from the St. Lawrence/Ottawa River populations and are a genetic intermediate that historically connected the southern (Mississippi River) and northern (St. Lawrence) populations across its range. Our study has shown that the Mississagi Hickorynut population is larger than previously known and

occurs from the river mouth to upstream of the town of Iron Bridge—a nearly six-fold increase in occupied reach. With the presence of multiple size classes, the observation of gravid females in the recent past (Zanatta and Woolnough 2011) and the continued persistence of its imperilled host (Pratt 2008; Bouvier *et al.* 2013; COSEWIC 2017), it is likely that a reassessment of this population's status could result in an overall assessment of fair to good (see Bouvier *et al.* 2013; DFO 2013).

The Mississagi River accounts for one of the largest inflows into Lake Huron and overall it is a large, well-oxygenated, clear system with a dominantly sandy substrate. Hickorynut is known to be found in large rivers, at deeper depths (>2 m) than other mussel species, and on small substrates, typically composed of sand (Clarke 1981; COSEWIC 2011). Based on the presence of the presumed host fish of this mussel and the information on depth and substrate collected at sites where Hickorynut was detected, the Mississagi River survey provides additional evidence to support the known preferred habitat of this Endangered species. Given that the habitat throughout the survey area appeared to be broadly consistent among sites, it is likely that there is much more available habitat and that there are more Hickorynut throughout the system.

While the Mississagi River and its freshwater mussel assemblage are isolated from many of the threats associated with urban and agricultural activities that have been implicated in mussel species declines across southwestern Ontario (COSEWIC 2013, 2021), the assemblage is not immune to risk. The construction and continued operation of the Red Rock Falls Generating Station represents a significant barrier to upstream movement of mussel hosts like Lake Sturgeon (Barth *et al.* 2018). The establishment of dreissenid mussels in the area beyond the delta similarly limits the downstream spread and connection with historical populations of freshwater mussels throughout the Great Lakes. Ongoing efforts to control the spread and impacts of invasive Sea Lamprey (*Petromyzon marinus*), an identified threat to the Hickorynut's presumed host, have highlighted the difficult intersection of invasive species management and species-at-risk protection (DFO 2021; Pratt *et al.* 2021). Newton *et al.* (2017) showed that granular Bayluscide® (Bayer CropScience, Inc., Calamba, Laguna, Philippines), a molluscicide commonly used to assess and treat larval Sea Lamprey populations in the Mississagi River, is highly toxic to many freshwater mussels and, to a lesser degree, young Lake Sturgeon (Boogaard *et al.* 2003). Andrews *et al.* (2021) reported that Hickorynut and Lake Sturgeon are each among the top five species with the highest potential

relative risk of mortality from Bayluscide when compared to other mussel and fish species of conservation concern. Modelling by Smyth and Drake (2021) indicates that although direct mortality is often low or zero during granular Bayluscide assessments (due to small application sizes and patchy mussel distributions), the potential for the deaths of hundreds of freshwater mussels and fish (particularly native lamprey species) following even a single assessment cycle remain. Bayluscide is also used in conjunction with TFM (3-trifluoromethyl-4-nitrophenol) during large scale, whole river treatments to eradicate larval Sea Lamprey in the Mississagi River. Bayluscide has been applied to the Mississagi River dozens of times since the 1960s (Daugherty *et al.* 1983; Andrews *et al.* 2021). The long-term implications of ongoing efforts to assess and control the impacts of invasive Sea Lamprey through Bayluscide application, in part designed to protect imperilled Lake Sturgeon with an indirect benefit to Hickorynut, are not fully understood. Understanding and mitigating threats such as this is essential to the long-term protection and restoration of the freshwater mussel assemblage within the Mississagi River as well as in other parts of Ontario.

The collaborative surveys conducted in the Mississagi River in 2023 provide a better understanding of freshwater mussel distributions in an understudied area and have specifically clarified the distribution of the Endangered Hickorynut. Future research in other rivers along the north shore of lakes Huron and Superior, where Lake Sturgeon occur, should be given priority as other undiscovered populations of Hickorynut may still occur. Bell (1861) reported mussel collections from other nearby rivers on the Canadian side of Lake Superior and Lake Huron watersheds including other well known Lake Sturgeon rivers (e.g., Goulais River). Although Bell did not find Hickorynut during his searches in these systems, more intensive surveys involving the application of SCUBA may prove fruitful. Currently, the Mississagi River represents the only known population of Hickorynut in Lake Huron and its occurrence and habitat should be protected. Future research on understanding Hickorynut population size structure, reproduction, and threat mitigation as well as the identification and protection of critical habitat will aid in the management and conservation of the species.

Author Contributions

Conceptualization: T.J.M., K.A.M.O., and K.S.; Funding Acquisition: T.J.M.; Investigation: K.A.M.O., T.J.M., and M.P.G.; Methodology: T.J.M., M.P.G., and K.A.M.O.; Writing – Original Draft: K.A.M.O. and T.J.M.; Writing – Review & Editing: K.A.M.O., T.J.M., M.P.G., and K.S.

Acknowledgements

We acknowledge that the work undertaken and reported here took place on the traditional and unceded territories of Indigenous Peoples of the Mississauga First Nation, part of the Mississauga Nation and a signatory to the Robinson Huron Treaty of 1850, that have lived in concert with the natural environment since time immemorial and will continue to do so into the future. We recognize the special bond that Indigenous Peoples have with the landscape and are thankful and grateful that our project took place on this beautiful landscape.

We thank all those involved in the field work for this project (Fisheries and Oceans Canada: J. Colm, L. Collinson, M. Vezina-Doré, S. Fitzgerald, S. Jacques, L. Savard, R. Leger Daigle, H.-F. Ellefsen; Environment and Climate Change Canada: B. Gray, C. Treen, C. Weaver, E. MacLennan-Nobrega; Mississauga First Nation Lands and Resources Department: R.L. Ferrigan, A. Chiblow, M. Moffatt, P. Morningstar). Thanks to all of those who provided support and input into planning (A. Price, D. Marson, K. Smith, D. Linley, M. Cowley, D. Campbell, W. Gardiner, J. Barnucz, L. O'Connor, T. Pratt, K. Bowen, W. Currie, and R. Rozon). Thanks to J. Colm for creation of maps and R. Dolson for statistical and R code support. Thanks to the Hickorynut distributional expertise of P. Badra, L. Kitchel, J. Trotter, D. Woolnough, and D. Zanatta. D.A. Price, Fisheries and Oceans Canada, Aquatic Invasive Species Program provided recent information on fish species present in the Mississagi River. The authors thank all of those who provide records used in the Lower Great Lakes Unionid Database. Thanks to the Hanner Laboratory at the University of Guelph who completed the molecular analysis and K. Layton for her editorial review of the methods section. Permits acquired for this work included *Species at Risk Act* (22-PCAA-00011) and a Letter of Authorization to Conduct Research in a Provincial Park or Conservation Reserve from Ontario Parks. Financial support for this project was provided by Fisheries and Oceans Canada's Species at Risk Program.

Literature Cited

- Aldridge, D.C., I.S. Ollard, Y.V. Bernalaya, I.N. Bolotov, K. Douda, J. Geist, W.R. Haag, M.W. Klunzinger, M. Lopes-Lima, M.C. Mlambo, N. Riccardi, R. Sousa, D.L. Stayer, S.H. Torres, C.C. Vaughn, T. Zajac, and A. Zieritz. 2022. Freshwater mussel conservation: a global horizon scan of emerging threats and opportunities. *Global Change Biology* 29: 575–589. <https://doi.org/10.1111/gcb.16510>
- Andrews, D.W., E.R.B. Smyth, D.E. Lebrun, T.J. Morris, K.A. McNichols-O'Rourke, and D.A.R. Drake. 2021. Relative risk of granular bayluscide applications for fishes and mussels of conservation concern in the Great Lakes basin. DFO Canadian Science Advisory Secretariat Research Document 2021/034. Accessed 17 January 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/40974339.pdf>.
- Barth, C.C., D. Burnett, C.A. McDougall, and P.A. Nelson. 2018. Information in support of the 2017 COSEWIC assessment and status report on the Lake Sturgeon (*Acipenser fulvescens*) in Canada. Canadian Manuscript Report of Fisheries and Aquatic Sciences 3166. Accessed 2 October 2024. https://publications.gc.ca/collections/collection_2019/mpo-dfo/Fs97-4-3166-eng.pdf.
- Bell, R. 1861. List of recent land and fresh-water shells collected around Lakes Superior and Huron in 1859–60. *Canadian Naturalist and Geologist, and the Proceedings of the Natural History Society* 6: 268–270. Accessed 30 January 2026. <https://www.biodiversitylibrary.org/page/7593465>.
- Boogaard, M.A., T.D. Bills, and D.A. Johnson. 2003. Acute toxicity of TFM and a TFM/Niclosamide mixture to selected species of fish, including Lake Sturgeon (*Acipenser fulvescens*) and Mudpuppies (*Necturus maculosus*), in laboratory and field exposures. *Journal of Great Lakes Research* 29: 529–541. [https://doi.org/10.1016/S0380-1330\(03\)70514-0](https://doi.org/10.1016/S0380-1330(03)70514-0)
- Bouvier, L.D., A. Paquet, and T.J. Morris. 2013. Information in support of a recovery potential assessment of Hickorynut (*Obovaria olivaria*) in Canada. DFO Canadian Science Advisory Secretariat Research Document 2013/041. Accessed 22 April 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/350232.pdf>.
- Brady, T., M. Hove, C. Nelson, R. Gordon, D. Hornback, and A. Kapuscinski. 2004. Suitable host fish species determined for hickorynut and pink heelsplitter. *Ellipsaria* 6: 14–15.
- Bucholz, J.R., N.M. Sard, N.M. VanTassel, J.D. Lozier, T.J. Morris, A. Paquet, and D.T. Zanatta. 2022. RAD-tag and mitochondrial DNA sequencing reveal the genetic structure of a widespread and regionally imperiled freshwater mussel, *Obovaria olivaria* (Bivalvia: Unionidae). *Ecology and Evolution* 12: e8560. <https://doi.org/10.1002/ece3.8560>
- Clarke, A.H. 1981. *The Freshwater Molluscs of Canada*. National Museum of Natural Sciences. National Museums of Canada, Ottawa, Ontario, Canada.
- Cliff, M., M. Hove, and M. Haas. 2001. Creeper glochidia appear to be host generalists. *Ellipsaria* 3: 18–19. Accessed 15 March 2024. https://molluskconservation.org/EVENTS/ELLIPSARIA/Ellipsaria2001_31.pdf.
- CNSC (Canadian Nuclear Safety Commission). 2023. Environmental protection review report: Blind River Refinery. Accessed 15 March 2024. <https://www.cnscc-ccsn.gc.ca/eng/resources/publications/reports/brr/>.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2011. COSEWIC assessment and status report on the Hickorynut *Obovaria olivaria* in Canada. COSEWIC, Ottawa, Ontario, Canada.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2013. COSEWIC status appraisal summary on the Kidneyshell *Ptychobranchus fasciolaris* in Canada. COSEWIC, Ottawa, Ontario, Canada.
- COSEWIC (Committee on the Status of Endangered

- Wildlife in Canada**). 2017. COSEWIC assessment and status report on the Lake Sturgeon *Acipenser fulvescens*, Western Hudson Bay populations, Saskatchewan-Nelson River populations, Southern Hudson Bay-James Bay populations and Great Lakes-Upper St. Lawrence populations in Canada. COSEWIC, Ottawa, Ontario, Canada.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada)**. 2021. COSEWIC assessment and status report on the Purple Wartyback *Cyclonaias tuberculata* in Canada. COSEWIC, Ottawa, Ontario, Canada.
- Daugherty, W.E., H.H. Moore, J.J. Tibbles, S.M. Dustin, and B.G.H. Johnson**. 1983. Sea Lamprey management in the Great Lakes. Annual Report for the year 1983 to Great Lakes Fishery Commission. Accessed 14 January 2026. https://www.glfco.org/pubs/slcp/annual_reports/ANN_UAL_REPORT_1983.pdf.
- DFO (Fisheries and Oceans Canada)**. 2013. Recovery Potential Assessment of Hickorynut (*Obovaria olivaria*) in Canada. DFO Canadian Science Advisory Secretariat Science Advisory Report 2013/041. Accessed 13 December 2023. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/349848.pdf>.
- DFO (Fisheries and Oceans Canada)**. 2020. Action plan for the Ausable River in Canada: an ecosystem approach. *Species at Risk Act* Action Plan Series. Fisheries and Oceans Canada, Ottawa, Ontario, Canada. Accessed 21 November 2023. https://wildlife-species.canada.ca/species-risk-registry/virtual_sara/files/plans/Ap-RiviereAusableRiver-v00-2020Mar-Eng.pdf.
- DFO (Fisheries and Oceans Canada)**. 2021. Science advice on the potential harm of granular Bayluscide applications to fish and mussel species at risk. DFO Canadian Science Advisory Secretariat Science Advisory Report 2021/016. Accessed 19 November 2025. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/4097604x.pdf>.
- Ferreira-Rodríguez, N., Y.B. Akiyama, O.V. Aksenova, R. Araujo, M.C. Barnhart, Y.V. Bepalaya, A.E. Bogan, I.N. Bolotov, P.B. Budha, C. Clavijo, S.J. Clearwater, G. Darrigran, V.T. Do, K. Douda, E. Froufe, C. Gumpinger, L. Henrikson, C.L. Humphrey, N.A. Johnson, O. Klishko, M.W. Klunzinger, S. Kovitvadhí, U. Kovitvadhí, J. Lajtner, M. Lopes-Lima, E.A. Moorkens, S. Nagayama, K.-O. Nagel, M. Nakano, J.N. Negishi, P. Ondina, P. Oulasvirta, V. Prié, N. Riccardi, M. Rudzite, F. Sheldon, R. Sousa, D.L. Strayer, M. Takeuchi, J. Taskinen, A. Teixeira, J.S. Tiemann, M. Urbańska, S. Varandas, M.V. Vinarski, B.J. Wicklow, T. Zając, and C.C. Vaughn**. 2019. Research priorities for freshwater mussel conservation assessment. *Biological Conservation* 231: 77–87. <https://doi.org/10.1016/j.biocon.2019.01.002>
- Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek**. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology Biotechnology* 3: 294–299.
- Gibson, M.P., K.A. McNichols-O'Rourke, and T.J. Morris**. 2024. Unionid monitoring and biodiversity observation (UMBO) network assessment in the Maitland River watershed, Ontario. Canadian Data Report of Fisheries and Aquatic Sciences 1395. Accessed 15 March 2024. https://publications.gc.ca/collections/collection_2024/mpo-dfo/Fs97-13-1395-eng.pdf.
- Government of Canada**. 2023. Species at Risk Public Registry: Species search - Molluscs. Accessed 7 February 2024. <https://species-registry.canada.ca/index-en.html#/species?taxonomyId=Molluscs&sortBy=commonNameSort&sortDirection=asc&pageSize=10>.
- Government of Ontario**. 2014. Species at Risk in Ontario (SARO) List. Ministry of the Environment, Conservation and Parks. King's Printer for Ontario, 2012–24. Accessed 15 February 2023. <https://www.ontario.ca/page/hickorynut>.
- Haag, W.R., and J.D. Williams**. 2014. Biodiversity on the brink: an assessment of conservation strategies for North American freshwater mussels. *Hydrobiologia* 735: 45–60. <https://doi.org/10.1007/s10750-013-1524-7>
- Harrison, M., V.A. Sotola, A. Zalmat, K.T. Sullivan, B.M. Littrell, T.H. Bonner, and N.H. Martin**. 2023. A comparison of non-destructive visceral swab and tissue biopsy sampling methods for genotyping-by-sequencing in the freshwater mussel *Fusconaia askewi*. *Genes (Basel)* 14: 1197. <https://doi.org/10.3390/genes14061197>
- Harvard University Museum of Comparative Zoology**. 2016. MCABASE: the database of the Zoological Collections. Accessed 10 April 2024. <http://mczbase.mcz.harvard.edu/guid/MCZ:Mala:269129>.
- Hatch**. 2010. Mississagi River Watershed Management Plan. Prepared for the Brookfield Renewable Power (BRP) and the Ontario Ministry of Natural Resources (MNR). Accessed 18 December 2023. https://evolugen.com/wp-content/uploads/2023/05/Volume-1-Main-Report-March-2010_Web.pdf.
- Hatch**. 2022. Flood risk assessment report for Huron Shores flood risk assessment. The Corporation of the Municipality of Huron Shores, Ontario Canada. Accessed 16 March 2024. https://huronshores.ca/wp-content/uploads/2022/09/H366743-0000-228-230-0003_Rev0-Flood-Risk-Assessment.pdf.
- Hayes Morris, C., and T.J. Morris**. 2024. The history of commercial freshwater mussel harvest in southern Ontario: a short-lived fishery with long-lasting consequences. *Canadian Field-Naturalist* 138: 205–216. <https://doi.org/10.22621/cfn.v138i3.3391>
- Henley, W.F., J.P. Grobler, and R.J. Neves**. 2006. Non-invasive method to obtain DNA from freshwater mussels (Bivalvia: Unionidae). *Journal of Shellfish Research* 25: 975–977. [https://doi.org/10.2983/0730-8000\(2006\)25\[975:nmtodf\]2.0.co;2](https://doi.org/10.2983/0730-8000(2006)25[975:nmtodf]2.0.co;2)
- Hoeh, W.R., and J.B. Burch**. 1989. The taxonomic status of *Anodonta lacustris* Lea (Bivalvia: Unionidae). *Walkerana* 3: 263–276.
- Keretz, S.S., D.A. Woolnough, T.J. Morris, E.F. Roseman, A.K. Elgin, and D.T. Zanatta**. 2021. Limited co-existence of native unionids and invasive dreissenid mussels more than 30 y post dreissenid invasion in a large river system. *American Midland Naturalist* 186: 157–175. <https://doi.org/10.1674/0003-0031-186.2.157>
- LGLUD (Lower Great Lakes Unionid Database)**. 2024. Lower Great Lakes Unionid Database. Microsoft Access 2016. Fisheries and Oceans Canada, Great Lakes Lab-

- oratory for Fisheries and Aquatic Sciences, Burlington, Ontario, Canada.
- Lopes-Lima, M., L.E. Burlakova, A.Y. Karatayev, K. Mehler, M. Seddon, and R. Sousa.** 2018. Conservation of freshwater bivalves at the global scale: diversity, threats and research needs. *Hydrobiologia* 810: 1–14. <https://doi.org/10.1007/s10750-017-3486710.1007/s10750-017-3486-7>
- Marson, D., E. Gertzen, and B. Cudmore.** 2016. Results of Fisheries and Oceans Canada's 2014 Asian Carp early detection field surveillance program. Canadian Manuscript Report Fisheries and Aquatic Sciences 3103. Accessed 2 October 2024. https://publications.gc.ca/collections/collection_2016/mpo-dfo/Fs97-4-3103-eng.pdf.
- Martel, A.L., H.F. Ellefsen, K. Iives, A. Paquet, N. Alfonso, M.H. Fraser, B. Aubrey, J. Heinerth, S. Jacques, M. D'Aguiar, L. Bassi, C. Rennie, and T. Haxton.** 2023. [Abstract] Research on the Hickorynut mussel–Lake Sturgeon interaction and their preferential habitats in the fluvial St. Lawrence River and its tributaries, including the Ottawa River. Pages 13–14 in *Proceedings of the 2023 Canadian Freshwater Mollusc Research Meeting: 15–16 November 2023*, Burlington, Ontario, Canada. Edited by T.J. Morris, M.P. Gibson, S.M. Reid, and K.A. McNichols-O'Rourke. Canadian Technical Report of Fisheries and Aquatic Sciences 3612. Accessed 13 April 2024. https://publications.gc.ca/collections/collection_2024/mpo-dfo/Fs97-6-3612-eng.pdf.
- Martel, A.L., and J.B. Madill.** 2017. [Poster] Host fish requirements help locate the rare Hickorynut mussel in the Ottawa River. Canadian Society of Zoologist Conference, 15–19 May 2017, Winnipeg, Manitoba, Canada. Accessed 15 March 2024. <https://www.ottawariverkeeper.ca/wp-content/uploads/2015/12/Obovaria-poster-CSZ-May-2017-12a.pdf>.
- Martel, A.L., and I. Picard.** 2005. The Hickorynut mussel, *Obovaria olivaria*, a deepwater unionid under scrutiny in Canadian rivers. *Tentacle* 13: 2–3.
- Martel, A.L., I. Picard, N. Binnie, B. Sawchuk, J. Madill, and F. Schueler.** 2006. The rare Olive Hickorynut mussel, *Obovaria olivaria*, in the Ottawa River, eastern Canada. *Tentacle* 14: 31–32.
- McNichols-O'Rourke, K.A., A. Robinson, and T.J. Morris.** 2012. Summary of freshwater mussel timed search surveys in southwestern Ontario in 2010 and 2011. Canadian Manuscript Report of Fisheries and Aquatic Sciences 3009. Accessed 14 March 2024. https://publications.gc.ca/collections/collection_2013/mpo-dfo/Fs97-4-3009-eng.pdf.
- MECP (Ministry of the Environment, Conservation and Parks).** 2021. Mississagi Delta Provincial Park Management Plan. Government of Ontario. Accessed 15 March 2024. <https://www.ontario.ca/page/mississagi-delta-provincial-park-management-plan>.
- MECP (Ministry of the Environment, Conservation and Parks).** 2024. Ontario Parks Park Locator. Government of Ontario. Accessed 14 March 2024. <https://www.ontario.parks.ca/park-locator>.
- Metcalfe-Smith, J.L., G.L. Mackie, J. Di Maio, and S.K. Staton.** 2000. Changes over time in the diversity and distribution of freshwater mussels (Unionidae) in the Grand River, southwestern Ontario. *Journal of Great Lakes Research* 26: 445–459. [https://doi.org/10.1016/S0380-1330\(00\)70707-6](https://doi.org/10.1016/S0380-1330(00)70707-6)
- Miller, E.N., K.A. Kidd, M.G. Surette, C.J. Bennett, J. Salerno, and P.L. Gillis.** 2022. Effects of municipal wastewater effluents on the digestive gland microbiome of wild freshwater mussels (*Lasmigona costata*). *Ecotoxicology and Environmental Safety* 241: 113774. <https://doi.org/10.1016/j.ecoenv.2022.113774>
- Miller, A.C., and B.S. Payne.** 1993. Qualitative versus quantitative sampling to evaluate population and community characteristics at a large-river mussel bed. *American Midland Naturalist* 130: 133–145. <https://doi.org/10.2307/2426282>
- MolluscaBase.** 2023. MolluscaBase. Accessed 15 December 2023. <https://www.molluscabase.org/>.
- MolluscaBase.** 2025. MolluscaBase. *Lampsilis radiata siliquioidea* (Barnes, 1823). Accessed 10 December 2025. <https://www.molluscabase.org/aphia.php?p=taxdetails&id=861844>.
- Morris, T.J., F. Gibson, K.A. McNichols-O'Rourke, M.N. Sheldon, and D.T. Zanatta.** 2020. [Abstract] Everything old is new again: the rediscovery of Lake Floater (*Pyganodon lacustris*) in Canada. Page 21 in *Proceedings of the 2019 Canadian Freshwater Mollusc Research Meeting: 3–4 December 2019*, Burlington, Ontario, Canada. Edited by T.J. Morris, K.A. McNichols-O'Rourke, and S.M. Reid. Canadian Technical Report of Fisheries and Aquatic Sciences 3352. Accessed 14 March 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/4084724x.pdf>.
- Morris, T.J., A.L. Martel, G. Wilson, M.N. Goguen, K.A. McNichols-O'Rourke, and J.E. Colm.** 2022. Guidance on the collection of freshwater mussel (Bivalvia: Unionida) vouchers within Canada. Canadian Manuscript Report of Fisheries and Aquatic Sciences 3241. Accessed 14 March 2024. https://publications.gc.ca/collections/collection_2022/mpo-dfo/Fs97-4-3241-eng.pdf.
- Morris, T.J., K.A. McNichols-O'Rourke, and A. Robinson.** 2012. A preliminary survey of the freshwater mussels of the Bayfield River watershed and nearby Lake Huron tributaries. Canadian Manuscript Report of Fisheries and Aquatic Sciences 2993. Accessed 2 October 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/349598.pdf>.
- NatureServe.** 2026. NatureServe Network Biodiversity Location Data accessed through NatureServe Explorer [web application]. NatureServe, Arlington, Virginia, USA. Accessed 29 January 2026. https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.113791/Obovaria_olivaria.
- Newton, T.J., M.A. Boogaard, B.R. Gray, T.D. Hubert, and N.A. Schloesser.** 2017. Lethal and sub-lethal responses of native freshwater mussels exposed to granular Bayluscide®, a Sea Lamprey larvicide. *Journal of Great Lakes Research* 43: 370–378. <https://doi.org/10.1016/j.jglr.2016.12.010>
- Nobles, T., and Y. Zhang.** 2011. Biodiversity loss in freshwater mussels: importance, threats, and solutions. Pages 137–162 in *Biodiversity Loss in a Changing Planet*. Edited by O. Grillo and G. Venora. InTch Publishing, Ri-

- jeka, Croatia. <https://doi.org/10.5772/25102>
- Obermeyer, B.K.** 1998. A comparison of quadrats versus timed snorkel searches for assessing freshwater mussels. *American Midland Naturalist* 139: 331–339. [https://doi.org/10.1674/0003-0031\(1998\)139\[0331:acoqvt\]2.0.co;2](https://doi.org/10.1674/0003-0031(1998)139[0331:acoqvt]2.0.co;2)
- Oksanen, J., G. Simpson, F. Blanchet, R. Kindt, P. Legendre, P. Minchin, R. O'Hara, P. Solymos, M. Stevens, E. Szoecs, H. Wagner, M. Barbour, M. Bedward, B. Bolker, D. Borcard, G. Carvalho, M. Chirico, M. De Caceres, S. Durand, H. Evangelista, R. FitzJohn, M. Friendly, B. Furneaux, G. Hannigan, M. Hill, L. Lahhti, D. McGlenn, M. Ouellette, E. Ribeiro Cunha, T. Smith, A. Stier, C. Ter Braak, and J. Weedon.** 2022. *vegan*: Community Ecology Package. R package version 2.6-4.
- Porto-Hannes, I., L.E. Burlakova, D.T. Zanatta, and H.R. Lasker.** 2021. Boundaries and hybridization in a secondary contact zone between freshwater mussel species (Family: Unionidae). *Heredity* 126: 955–973. <https://doi.org/10.1038/s41437-021-00424-x>
- Pratt, T.C.** 2008. Population status and threats of Lake Sturgeon in Designatable Unit 8 (Great Lakes / St. Lawrence River Watersheds). DFO Canadian Science Advisory Secretariat Research Document 2008/043. Accessed 16 October 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/334960.pdf>.
- Pratt, T.C., B.J. Morrison, H.R. Quinlan, R.F. Elliott, S.A. Grunder, J.A. Chiotti, and B.A. Young.** 2021. Implications of the Sea Lamprey control program for Lake Sturgeon conservation and rehabilitation efforts. *Journal of Great Lakes Research* 41: S421–S429. <https://doi.org/10.1016/j.jglr.2020.06.014>
- R Core Team.** 2023. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ratnasingham, S., and P.D.N. Hebert.** 2007. BOLD: the Barcode of Life Data System. *Molecular Ecology Notes* 7: 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Reid, S.M., and T.J. Morris.** 2017. Tracking the recovery of freshwater mussel diversity in Ontario rivers: evaluation of a quadrat-based monitoring protocol. *Diversity* 9: 1–17. <https://doi.org/10.3390/d9010005>
- SARA (Species at Risk Act) Public Registry.** 2024a. Species summary, Hickorynut (*Obovaria olivaria*). Government of Canada, Ottawa, Ontario, Canada. Accessed 27 February 2025. <https://species-registry.canada.ca/index-en.html#/species/1150-806>.
- SARA (Species at Risk Act) Public Registry.** 2024b. Species summary, Lake Sturgeon (*Acipenser fulvescens*), Great Lakes-Upper St. Lawrence populations. Government of Canada, Ottawa, Ontario, Canada. Accessed 14 March 2024. <https://species-registry.canada.ca/index-en.html#/species/842-601>.
- Schloesser, D.W., J.L. Metcalfe-Smith, W.P. Kovalak, G.D. Longton, and R.D. Smithee.** 2006. Extirpation of freshwater mussels (Bivalvia: Unionidae) following the invasion of dreissenid mussels in an interconnecting river of the Laurentian Great Lakes. *American Midland Naturalist* 155: 307–320. [https://doi.org/10.1674/0003-0031\(2006\)155\[307:eofimbu\]2.0.co;2](https://doi.org/10.1674/0003-0031(2006)155[307:eofimbu]2.0.co;2)
- Smyth, E.R.B., and D.A.R. Drake.** 2021. Estimating the mortality of fishes and mussels of conservation concern resulting from Bayluscide applications within four rivers of the Huron-Erie Corridor. DFO Canadian Science Advisory Secretariat Research Document 2021/035. Accessed 02 October 2024. <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/40974595.pdf>.
- Stanfield, L.** 2010. Ontario stream assessment protocol. Version 8. Fisheries Policy Section. Ontario Ministry of Natural Resources. Peterborough, Ontario, Canada. Accessed 14 February 2022. <https://docs.ontario.ca/documents/2575/226871.pdf>.
- Vaughn, C.C.** 2018. Ecosystem services provided by freshwater mussels. *Hydrobiologia* 810: 15–27. <https://doi.org/10.1007/s10750-017-3139-x>
- Vaughn, C.C., and T.J. Hoellein.** 2018. Bivalve impacts in freshwater and marine ecosystems. *Annual Review of Ecology, Evolution, and Systematics* 49: 183–208. <https://doi.org/10.1146/annurev-ecolsys-110617-062703>
- Zanatta, D.T., and D.A. Woolnough.** 2011. Confirmation of *Obovaria olivaria*, Hickorynut mussel (Bivalvia: Unionidae), in the Mississagi River, Ontario, Canada. *Northeastern Naturalist* 18: 1–6. <https://doi.org/10.1656/045.018.0101>

Received 20 November 2024

Accepted 30 January 2026

Associate Editor: D.A.W. Lepitzki

SUPPLEMENTARY MATERIALS:

FIGURES S1–S9. Digital voucher specimens.

FIGURE S1. Eastern Elliptio (*Elliptio complanata*).

FIGURE S2. Plain Pocketbook (*Lampsilis cardium*).

FIGURE S3. Fatmucket (*Lampsilis siliquoidea*).

FIGURE S4. Hickorynut (*Obovaria olivaria*).

FIGURE S5. Giant Floater (*Pyganodon grandis*).

FIGURE S6. Lake Floater (*Pyganodon lacustris*).

FIGURE S7. Creeper (*Strophitus undulatus*).

FIGURE S8. Flutedshell (*Lasmigona costata*) shell.

FIGURE S9. Black Sandshell (*Ligumia recta*) shell.

FIGURE S10. Habitat variables at sites with (present) and without (absent) Hickorynut (*Obovaria olivaria*) in the Mississagi River, Ontario in 2023.

TABLE S1. Information collected at each site surveyed in 2023 on the Mississagi River, Ontario.